

THE CONCEPT OF BOUNDED MEAN MOTION FOR TORAL HOMEOMORPHISMS

T. Jäger*

April 30, 2008

Abstract

A conservative irrational pseudo-rotation of the two-torus is semi-conjugate to the irrational rotation if and only if it has the property of *bounded mean motion* [1]. (Here ‘irrational pseudo-rotation’ means a toral homeomorphism with a unique and totally irrational rotation vector.) The aim of this note is to explore this concept some further. For instance, we provide an example which shows that the preceding statement does not hold in the non-conservative case. Further, we collect a number of observations concerning the case where the bounded mean motion property fails. In particular, we show that a non-wandering irrational pseudo-rotation of the two-torus with unbounded mean motion has sensitive dependence on initial conditions.

1 Introduction

We denote by $\text{Homeo}_0(\mathbb{T}^d)$ the set of homeomorphisms of the d -dimensional torus which are homotopic to the identity. An important topological and dynamical invariant for this class of maps is given by the rotation set. Given $f \in \text{Homeo}_0(\mathbb{T}^d)$ with lift $F : \mathbb{R}^d \rightarrow \mathbb{R}^d$, it is defined as

$$(1.1) \quad \rho(F) := \left\{ \rho \in \mathbb{R}^d \mid \exists n_i \nearrow \infty, z_i \in \mathbb{R}^d : \lim_{i \rightarrow \infty} (F^{n_i}(z_i) - z_i)/n_i = \rho \right\}.$$

The set $\rho(F)$ is always compact, and if $d = 2$ it also convex [2]. At least in dimension two, it is well-known that the shape of $\rho(F)$ is closely related to the dynamical properties of f . For instance, if $\rho(F)$ has non-empty interior, then all rational rotation vectors in $\text{int}(\rho(F))$ are realised by periodic orbits [3], the topological entropy of f is strictly positive [4] and the rotation set is continuous in f with respect to the topology of uniform convergence [5] (in general, the dependence on f is only upper semi-continuous).

In contrast to this, the situation is less well-understood when the rotation set is ‘thin’, meaning that it has empty interior. A particular case is that of an (*irrational*) *pseudo-rotation*, which is defined as a toral homeomorphism f whose rotation set is reduced to a single (totally irrational) rotation vector ρ . In this situation, we say f has *bounded mean motion*, if the *deviations from the constant rotation*

$$(1.2) \quad D(n, z) := F^n(z) - z - n\rho$$

are uniformly bounded in θ, x and n . This notion already played an important role in the description of the dynamics of certain skew product transformations, like quasiperiodically forced circle homeomorphisms [6, 7, 8] and almost periodically forced circle flows [9, 10]. The fact that it is equally important for understanding, and classifying, the possible dynamics of irrational pseudo-rotations is documented by the following statement.

Theorem 1.1 ([1]). *Suppose $f \in \text{Homeo}_0(\mathbb{T}^d)$ is minimal, or $d = 2$ and f is conservative. Then f is semi-conjugate to an irrational rotation on the d -dimensional torus if and only if it is an irrational pseudo-rotation with bounded mean motion.*

*Collège de France, Paris. Email: tobias.jager@college-de-france.fr

At least of the minimal/conservative case, this result suggests to divide irrational pseudo-rotations into two classes, those with and those without bounded mean motion. However, the usefulness of such a dichotomy will strongly depend on the information which can be deduced from the unboundedness of mean motion. Furthermore, it would certainly be desirable to weaken the recurrence assumption if possible.

In other words, Theorem 1.1 raises two obvious questions: (i) What happens in the non-minimal/dissipative case? (ii) What happens if the bounded mean motion property fails? While being far from giving a complete picture, the aim of this note is to collect a number of results and observations related to these two questions.

Concerning the first, we will provide examples of irrational pseudo-rotations with bounded mean motion which are not semi-conjugate to an irrational rotation (see Section 2). These examples have a very simple structure, namely they are skew-products over Denjoy counterexamples, with rotations on the fibres. Hence, they are still semi-conjugate to a one-dimensional irrational rotation, and they have very bad recurrence properties, since they admit homotopically non-trivial wandering open sets. We have to leave open here whether more sophisticated examples without these two properties exist. In particular, it is still possible that the assumptions in Theorem 1.1 could be weakened to some extent, for example to the non-existence of wandering open sets.

In order to address the second question, we will first of all consider the behaviour of the quantities $D_n(\theta, x)$ themselves in the case of unbounded mean motion. This is motivated by the fact that a good understanding of this behaviour already turned out to be crucial in the theory of the skew product transformations mentioned above. However, in order to obtain a good picture, some recurrence assumption on the system is needed. The best description can be given for minimal systems (see Section 3). Skew-products over Denjoy counterexamples allow again to demonstrate that a number of natural properties that hold in the minimal case cannot be expected in general.

Finally, in Section 4 we show that for an irrational pseudo-rotation of the two-torus without wandering open sets, the unboundedness of mean motion implies sensitive dependence on initial conditions. The interest of this result lies in particular in the fact that it demonstrates how the unboundedness of mean motion allows to obtain further information about the dynamics of the system.

Notation and terminology. For the sake of readability, we will use a number of conventions which might not be absolutely consistent from a strictly formal point of view, but should not cause any ambiguities and greatly simplify notation. For example, we will identify points in $\mathbb{T}^d$ with their lifts whenever the particular choice of the lift does not matter. Whenever there is a canonical distance function on a metric space, we will denote it by d . In the case of $\mathbb{R}$, $\mathbb{R}^d$ or $\mathbb{T}^d$, this will always be the usual Euclidean distance. The ε -neighbourhood of a point z in a metric space X will be denoted by $B_\varepsilon(z) = \{x \in X \mid d(z, x) < \varepsilon\}$. The Euclidean length of a vector $v \in \mathbb{R}^d$ will simply be denoted by $\|v\|$. Given a vector $v \in \mathbb{R}^d$ and a subset $S \subseteq \mathbb{R}^d$, we let $S + v := \{s + v \mid s \in S\}$.

Quotient maps like $\mathbb{R} \rightarrow \mathbb{T}^1$ or $\mathbb{R}^d \rightarrow \mathbb{T}^d$ will all be denoted by π , in product spaces π_i will be the projection to the respective coordinate. The rotation on $\mathbb{T}^d$ by an element $\rho \in \mathbb{T}^d$ will be denoted by R_ρ , as well as the translation by a vector $\rho \in \mathbb{R}^d$ on $\mathbb{R}^d$.

Acknowledgements. I would like to thank Sylvain Crovisier and Frédéric Le Roux for stimulating discussions. Further, I would like to thank J.-C. Yoccoz and the Collège de France for their hospitality and support during a long-term visit, and acknowledge support from the German Science Foundation (research fellowship Ja 1721/1-1).

2 Counter-examples to the existence of a semi-conjugacy

In this section, we provide examples of irrational pseudo-rotations which have bounded mean motion, but are not semi-conjugate to an irrational rotation. The main idea is to use skew products over Denjoy counter-examples, with rotations on the fibres. The dynamics

over the wandering intervals can then be modified, without affecting the rotation set, in order to produce some irregular behaviour. Since we will employ this basic construction several times, we first want to give a brief outline.

Given a rotation vector $\rho = (\rho_1, \rho_2) \in (\mathbb{R} \setminus \mathbb{Q}) \times \mathbb{R}$, we choose an orientation-preserving circle homeomorphism g with rotation number $\rho(g) = \rho_1$ and unique minimal set $M \subsetneq \mathbb{T}^1$. Further, we choose some continuous function $\beta : \mathbb{T}^1 \rightarrow \mathbb{R}$ which satisfies $\beta(x) = 0 \ \forall x \in M$ and define

$$(2.1) \quad f : \mathbb{T}^1 \times \mathbb{R} \rightarrow \mathbb{T}^1 \times \mathbb{R} \quad , \quad (x, y) \mapsto (g(x), y + \rho_2 + \beta(x) \bmod 1) .$$

This yields an irrational pseudo-rotation f with unique rotation vector ρ . One way to see this is to note that any f -invariant measure μ must project down to the unique g -invariant measure ν . Hence, the topological support of μ is contained in the set $M \times \mathbb{T}^1$. On this set, the replacement function $D(1, z) = F(z) - z$ (defined via a suitable lift F of f) is constant and equal to ρ , and therefore $\int_{\mathbb{T}^2} F(z) - z \, d\mu = \rho$ for all f -invariant measures μ . This implies $\rho(F) = \{\rho\}$ as claimed. The important point is that on the set M^c the function β can still be chosen arbitrarily in order to produce the desired phenomena.

As a first application of this construction, we prove the following:

Proposition 2.1. *Given any totally irrational rotation vector $\rho = (\rho_1, \dots, \rho_d) \in \mathbb{R}^d$, $d \geq 2$, there exists an irrational pseudo-rotation $f \in \text{Homeo}_0(\mathbb{T}^d)$ which has rotation vector ρ and bounded mean motion, but which is not semi-conjugate to the irrational rotation R_ρ .*

We will actually carry out the construction only in dimension $d = 2$, but the modifications for the general case are minor. Before we turn to the proof, we need the following lemma.

Lemma 2.2. *Let $\rho = (\rho_1, \rho_2) \in \mathbb{R}^2$ be totally irrational. Suppose $g \in \text{Homeo}_0(\mathbb{T}^1)$ is semi-conjugate to $R_{\rho_1} : \mathbb{T}^1 \rightarrow \mathbb{T}^1$ by a semi-conjugacy ϕ and $f \in \text{Homeo}_0(\mathbb{T}^2)$ is semi-conjugate to $R_\rho : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ by a semi-conjugacy ψ . Further, assume that f is a skew-product over g , that is $\pi_1 \circ f(x, y) = g(x) \ \forall (x, y) \in \mathbb{T}^2$.*

- (a) *If f has a unique minimal set, then the semi-conjugacy ψ can be chosen such that $\pi_1 \circ \psi(x, y) = \phi(x) \ \forall (x, y) \in \mathbb{T}^2$.*
- (b) *If f is given by (2.1), with $\beta|_M = 0$, then ψ can be chosen such that $\psi(x, y) = (\phi(x), y) \ \forall (x, y) \in M \times \mathbb{T}^1$.*

Proof. The argument is a slight variation of the one in [11, Section 4.14]. Choose lifts G, Φ, F and Ψ of the maps g, ϕ, f and ψ to $\mathbb{R}$ and $\mathbb{R}^2$, respectively, such that $\Phi \circ G = R_{\rho_1} \circ \Phi$, $\Psi \circ F = R_\rho \circ \Psi$ and $\pi_1 \circ F(x, y) = G(x)$. We claim that the function $\eta = \pi_1 \circ \Psi - \Phi \circ \pi_1$ is constant. In order to see this, note first that

$$\begin{aligned} \eta \circ F(x, y) &= \pi_1 \circ \Psi \circ F(x, y) - \Phi \circ \pi_1 \circ F(x, y) \\ &= \pi_1 \circ R_\rho \circ \Psi(x, y) - \Phi \circ G(x, y) \\ &= R_{\rho_1} \circ \pi_1 \circ \Psi(x, y) - R_{\rho_1} \circ \Phi(x) \\ &= \pi_1 \circ \Psi(x, y) - \Phi(x) = \eta(x, y) . \end{aligned}$$

Hence η is F invariant, and we can interpret it as a f -invariant function $\mathbb{T}^2 \rightarrow \mathbb{R}$. Since η is continuous, it follows first that it must be constant on the unique minimal set of f , and since any point must contain this minimal set in its ω -limit, η must be constant everywhere. Composing Ψ with a rotation in the x -direction, we can assume that $\eta = 0$. This proves part (a).

Now assume that f is of the form (2.1). Let $\mu = \text{Leb}_{\mathbb{T}^1} \circ \phi$. Then μ is g -invariant, and $\mu \times \text{Leb}_{\mathbb{T}^1}$ is f -invariant. Furthermore, ψ has to map $\mu \times \text{Leb}_{\mathbb{T}^1}$ to the unique R_ρ -invariant measure $\text{Leb}_{\mathbb{T}^2}$. This means that for μ -a.e. x , the map $\psi_x : y \mapsto \pi_2 \circ \psi(x, y)$ must preserve Lebesgue measure and hence be a rotation. By continuity, this is true for all $x \in \text{supp}(\mu) = M$. Using the notation $f_x(y) = \pi_2 \circ f(x, y)$, this further implies that for all $x \in M$ there holds

$$\begin{aligned} \psi_{g(x)}(y) + \rho_2 &= \psi_{g(x)}(y + \rho_2) = \psi_{g(x)} \circ f_x(y) \\ &= R_{\rho_2} \circ \psi_x(y) = \psi_x(y) + \rho_2 . \end{aligned}$$

Thus the rotations ψ_x and $\psi_{g(x)}$ are the same, and using continuity of ψ and minimality of M once more we obtain that $x \mapsto \psi_x$ is constant on M . Composing ψ with a rotation in the y -direction, we may therefore assume that $\psi_x = \text{Id}_{\mathbb{T}^1} \forall x \in M$, which proves statement (b). $\square$

Proof of Proposition 2.1. As mentioned, we give the proof in dimension 2. Let $\rho = (\rho_1, \rho_2)$ be totally irrational and define f as in (2.1), where we choose β as follows. First, let $(\alpha_n)_{n \in \mathbb{N}}$ be a sequence of real numbers, such that

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$;
- (ii) $\sup_{n \in \mathbb{Z}} |\sum_{i=0}^n \alpha_i| \leq 1$;
- (iii) $\limsup_{n \rightarrow \infty} \sum_{i=0}^n \alpha_i = 1$;
- (iv) $\liminf_{n \rightarrow \infty} \sum_{i=0}^n \alpha_i = 0$.

Let $I_0 = [a_0, b_0]$ be a connected component of M^c and let $I_n = [a_n, b_n] = g^n(I_0)$. Further, fix some $x_0 \in (a_0, b_0)$ and let $x_n = g^n(x_0)$. Choose a continuous function $\beta_0 : I_0 \rightarrow [0, 1]$ which satisfies $\beta_0(a_0) = \beta_0(b_0) = 0$ and $\beta_0(x_0) = 1$. Then define

$$\beta(x) := \begin{cases} \alpha_n \beta_0(g^{-n}(x)) & \text{if } x \in I_n \text{ for some } n \in \mathbb{Z} \\ 0 & \text{otherwise} \end{cases}.$$

Property (ii) of the α_n then implies that f has bounded mean motion. We claim that f cannot be semi-conjugate to the irrational rotation R_ρ . Suppose for a contradiction that ψ is a semi-conjugacy between f and R_ρ . By Lemma 2.2, we can assume that ψ is of the form $\psi(x, y) = (\phi(x), \psi_x(y))$, where ϕ is a semi-conjugacy between g and R_{ρ_1} , and $\psi_x = \text{Id}_{\mathbb{T}^1} \forall x \in M$. Due to uniform continuity, there exists $\delta > 0$, such that for all $z, z' \in \mathbb{T}^2$ with $d(z, z') < \delta$ there holds $d(\psi(z), \psi(z')) < \frac{1}{8}$. If we choose $k \in \mathbb{N}$ such that $d(x_k, b_k) < \delta$ (note that $|I_k| \rightarrow 0$ as $k \rightarrow \infty$), then it follows that for all $n \in \mathbb{Z}$ there holds

$$\begin{aligned} (2.2) \quad & d(\psi \circ f^n(x_k, 0), (\phi(b_{k+n}), n\rho_2)) = \\ & = d(\psi \circ f^n(x_k, 0), \psi \circ f^n(b_k, 0)) \\ & = d(R_\rho^n \circ \psi(x_k, 0), R_\rho^n \circ \psi(b_k, 0)) < 1/8. \end{aligned}$$

On the other hand, there holds $\pi_2 \circ f^n(x_k, 0) = \sum_{i=k}^{k+n-1} \alpha_i \bmod 1$. Hence, due to properties (i), (iii) and (iv) above there exist infinitely many $n \in \mathbb{N}$, such that

$$d(\pi_2 \circ f^n(x_k, 0), n\rho_2 + 1/2) < \delta/2.$$

If we choose such an n with the additional property that $d(x_{k+n}, b_{k+n}) < \frac{\delta}{2}$, then

$$d(f^n(x_k, 0), (b_{k+n}, n\rho_2 + 1/2)) < \delta.$$

Since $\psi(b_{k+n}, n\rho_2 + 1/2) = (\phi(b_{k+n}), n\rho_2 + 1/2)$, it follows that

$$(2.3) \quad d(\psi \circ f^n(x_k, 0), (\phi(b_{k+n}), n\rho_2 + 1/2)) < 1/8.$$

This clearly contradicts (2.2). $\square$

3 Unbounded Mean Motion

We first want to mention that the existence of irrational pseudo-rotations with unbounded mean motion is well-known. In [11], Herman provides examples of homeomorphisms of the two-torus which are skew products over an irrational rotation (so-called quasiperiodically forced circle homeomorphisms) and have totally irrational rotation vector, but are not semi-conjugate to the corresponding irrational rotation. Since for such skew products the statement of Theorem 1.1 holds without any recurrence assumption [7], these examples have unbounded mean motion. Other examples of irrational pseudo-rotations with unbounded

mean motion are provided by uniquely ergodic and weakly mixing torus diffeomorphisms, as constructed for example in [12].

In order to give a more detailed discussion, we first need a refined version of the notions introduced in Section 1. Given $f \in \text{Homeo}_0(\mathbb{T}^d)$ with lift F and a subset $A \subseteq \mathbb{T}^d$, we define the rotation set of f on A as

$$(3.1) \quad \rho_A(F) := \left\{ \rho \in \mathbb{R}^d \mid \exists n_i \nearrow \infty, x_i \in A : \lim_{i \rightarrow \infty} (F^{n_i}(x_i) - x_i)/n_i = \rho \right\}.$$

Further, when v is a vector in $\mathbb{R}^d \setminus \{0\}$, $\lambda \in \mathbb{R}$ and $\rho_A(F) \subseteq \lambda v + \{v\}^\perp$, then for any $z \in A$ we define the *deviations from the constant rotation parallel to v* as

$$(3.2) \quad D_v(n, z) := \langle F^n(z) - z - n\rho, v \rangle.$$

Here ρ can be any vector in $\lambda v + \{v\}^\perp$. When the quantities $D_v(n, z)$ are uniformly bounded, we say f has *bounded deviations parallel to v on A* .

A priori, it is not entirely clear whether there exist irrational pseudo-rotations with unbounded mean motion in all directions (that is, parallel to all vectors $v \in \mathbb{R}^d \setminus \{0\}$). For the examples mentioned above, there always exists some vector v , such that the deviations parallel to v are bounded. Hence, for the sake of completeness, we will provide examples with unbounded deviations parallel to all vectors at the end of this section (see Proposition 3.11).

Before, we want to gather some more information about the behaviour of the quantities $D_v(n, z)$ when the mean motion parallel to v is not bounded. In spirit, the results we present below are close to those collected in [6] for quasiperiodically forced circle homeomorphisms.

In the whole section, we suppose that $f \in \text{Homeo}_0(\mathbb{T}^d)$, F is a lift of f and $K \subseteq \mathbb{T}^d$ is a compact f -invariant subset. Further, we always assume that λ is a real number and v is a vector in $\mathbb{R}^d \setminus \{0\}$. As a first observation, we note that there always exists orbits on which the deviations are semi-bounded in the following sense.

Lemma 3.1. *Suppose that $\rho_K(F) \subseteq \lambda v + \{v\}$. Then given any $\varepsilon > 0$, there exist $z_-, z_+ \in K$, such that*

$$\sup_{n \in \mathbb{N}} D_v(n, z_-) < \varepsilon \quad \text{and} \quad \inf_{n \in \mathbb{N}} D_v(n, z_+) > -\varepsilon.$$

Proof. Without loss of generality, we may assume $\|v\| = 1$. Suppose for a contradiction that for all $z \in K$ there holds

$$\sup_{n \in \mathbb{N}} D_v(n, z) \geq \varepsilon > 0.$$

Then for any $z \in K$, there exists an integer $n(z)$ and a small neighbourhood $U(z)$, such that

$$\langle F^{n(z)}(z') - z', v \rangle - n(z)\lambda \geq \varepsilon/2 \quad \forall z' \in U(z).$$

Using compactness, we can find $z_1, \dots, z_m$ with $K \subseteq \bigcup_{i=1}^m U(z_i)$. Let $M := \max_{i=1}^m n(z_i)$. Inductively define a sequence of integers N_j and points ζ_j as follows: Let $N_0 = 0$ and ζ_0 be arbitrary. If $N_0, \dots, N_j$ and $\zeta_0, \dots, \zeta_j$ are given, choose $k \in \{1, \dots, m\}$ with $\zeta_j \in U(z_k)$ and let $\zeta_{j+1} := F^{n(z_k)}(\zeta_j)$ and $N_{j+1} := N_j + n(z_k)$. Then

$$\begin{aligned} \langle (F^{N_j}(z_0) - z_0), v \rangle &= \langle z_j - z_0, v \rangle \\ &= \sum_{i=0}^{j-1} \langle z_{i+1} - z_i, v \rangle \geq N_j \lambda + (j\varepsilon)/2 \geq (\lambda + \varepsilon/2M) \cdot N_j. \end{aligned}$$

By going over to a subsequence if necessary, we may further assume that $(F^{N_j}(z_0) - z_0)/N_j \rightarrow \rho \in \rho(F)$ as $j \rightarrow \infty$. Thus we obtain

$$\langle \rho, v \rangle = \lim_{j \rightarrow \infty} \langle (F^{N_j}(z_0) - z_0)/N_j, v \rangle \geq \lambda + \varepsilon/2M,$$

contradicting the assumptions. $\square$

In general, it seems difficult to say much more about the qualitative behaviour of the quantities $D_v(n, z)$. For instance, even when the deviations from the constant rotation are not uniformly bounded, they may be bounded on every single orbit, as the following example shows.

Example 3.2. (a) Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$ is a skew product over a Denjoy counter-example g , of the same form as in (2.1). Let $I = (a, b)$ be a connected component of M^c , where M is the unique minimal set of g , and $I_n = g^n(I)$. In order to define β in (2.1), choose a continuous function $\beta_0 : [a, b] \rightarrow [0, 1]$ with $\beta_0(a) = \beta_0(b) = 0$ and $\beta(x_0) = 1$ for some $x_0 \in I$. Let

$$\beta(x) := \begin{cases} \beta_0 \circ g^{-n}(x)/|n| & \text{if } x \in I_n \text{ with } n \leq 0 \\ 0 & \text{otherwise} \end{cases}.$$

Then it is easy to see that f does not admit bounded mean motion, but $\sup_{n \in \mathbb{N}} \|D(n, z)\| < \infty \forall z \in \mathbb{T}^2$.

(b) Evidently, the preceding example still exhibits orbits with unbounded mean motion backwards in time, that is $\sup_{n \in \mathbb{N}} \|D(-n, z)\| = \infty$ for some z . However, even this need not be true, as a slight modification of the construction shows. Suppose I^k , $k \in \mathbb{N}$ is a sequence of disjoint open subintervals of I and let $I_n^k := g^n(I^k)$. Choose $\beta_0 : I \rightarrow [0, 1]$, such that $\sup_{x \in I^k} \beta_0(x) = 1/k$. Then define

$$\beta(x) := \begin{cases} \beta_0 \circ g^n(x)/n & \text{if } x \in I_n^k \text{ with } |n| \leq 2^{k^2} \\ 0 & \text{otherwise} \end{cases}.$$

Since $\sum_{n=1}^{2^{k^2}} 1/n \geq k^2$, we have $\sup_{n \in \mathbb{N}} \|D(n, (x, 0))\| \geq k$ for some $x \in I^k$, and therefore $\sup_{n \in \mathbb{N}, z \in \mathbb{T}^2} \|D(n, z)\| = \infty$, but on the other hand there holds $\sup_{n \in \mathbb{Z}} \|D(n, z)\| < \infty \forall z \in \mathbb{T}^2$.

(c) A slight modification of these constructions also shows that the deviations from the constant direction can be unbounded in one direction, but at the same time bounded in the opposite direction. More precisely, it is possible to have $\sup_{n \in \mathbb{N}} D_v(n, z) = \infty$ for some $z \in \mathbb{T}^d$, but $\inf_{n \in \mathbb{Z}, z \in \mathbb{T}^d} D_v(n, z) > -\infty$, and vice versa.

The situation becomes different when f has some topological recurrence properties. Starting point is the following simple observation. Let $\mathcal{O}^+(z) = \{f^n(z) \mid n \in \mathbb{N}\}$ denote the forward orbit of z .

Lemma 3.3. Suppose $\rho(F) \subseteq \lambda v + \{v\}^\perp$ and $\sup_{n \in \mathbb{N}} |D_v(n, z)| \leq C < \infty$ for some $z \in \mathbb{T}^d$. Then $\sup_{n \in \mathbb{N}} |D_v(n, z)| \leq 2C \forall z \in \overline{\mathcal{O}^+(z)}$.

Proof. Suppose $D_v(n, z_0) > 2C$ for some $z_0 \in \overline{\mathcal{O}^+(z)}$. Then for some sufficiently small neighbourhood U of z_0 , there holds $D_v(n, z') > 2C \forall z' \in U$. In particular, when $f^k(z) \in U$, then $D_v(n, f^k(z)) > 2C$. However, this contradicts

$$D_v(n, f^k(z)) = D_v(n+k, z) - D_v(k, z) \leq 2C.$$

$\square$

Now suppose $f|_K$ is topologically transitive and denote the set of all points in K with dense forward orbit by $\mathcal{T}_K$. Recall that $\mathcal{T}_K$ is a residual (meaning dense G_δ) subset of K .

Corollary 3.4. Suppose $f|_K$ is topologically transitive and $\rho_K(F) \subseteq \lambda v + \{v\}^\perp$. If there holds $\sup_{n \in \mathbb{N}, z \in K} |D_v(n, z)| = \infty$, then $\sup_{n \in \mathbb{N}} |D_v(n, z)| = \infty \forall z \in \mathcal{T}_K$.

This statement is particularly interesting, when $f|_K$ is minimal, since this means that $\mathcal{T}_K = K$ and therefore all orbits have unbounded deviations. Together with Lemma 3.1, this entails the following corollary.

Corollary 3.5. *Suppose f is minimal, $\rho_K(F) \subseteq \lambda v + \{v\}^\perp$ and the deviations from the constant rotation parallel to v are not uniformly bounded on K . Then $f|_K$ has sensitive dependence on initial conditions.*

Proof. Due to uniform continuity, there exists $\varepsilon > 0$, such that $d(z_1, z_2) < \varepsilon$ implies $d(F(z_1), F(z_2)) < 1/2$. Suppose $\delta > 0$ and $z \in \mathbb{T}^d$ are given. We have to show that there exist $z_1, z_2 \in B_\delta(z)$ and $n \in \mathbb{N}$, such that $d(f^n(z_1), f^n(z_2)) \geq \varepsilon$. In order to do so, choose $z_1, z_2 \in B_\delta(z)$ with the properties that $\sup_{n \in \mathbb{N}} D_v(n, z_1) < \infty$ and $\inf_{n \in \mathbb{N}} D_v(n, z_2) > -\infty$. This is possible due to Lemma 3.1, minimality, and the fact that these properties are invariant under iteration by f .

Since the deviations are unbounded on all orbits by Lemma 3.4, the distance between $F^n(z_1)$ and $F^n(z_2)$ in $\mathbb{R}^d$ must become arbitrarily large. The choice of ε above therefore implies that for some $n \in \mathbb{N}$ there holds $d(F^n(z_1), F^n(z_2)) \in [\varepsilon, 1/2)$. However, as long as two points have distance less than $1/2$ in $\mathbb{R}^d$, their projections have exactly the same distance in $\mathbb{T}^d$. Hence $d(f^n(z_1), f^n(z_2)) \geq \varepsilon$ as required. $\square$

The preceding statement provides a first indication of how information on the behaviour of the deviations $D(n, z)$ may be used in order to draw further conclusions about the dynamical properties of f . In the proof, we have used that on the one hand the vectors $D(n, z)$ become arbitrarily large, but on the other hand we have also exploited the fact that they grow in different directions, depending on the starting point, due to Lemma 3.1.

In other words, we have not only made use of information on the size, but also on the behaviour of the directions of the vectors $D(n, z)$. In the following, we want to obtain a more detailed description of this behaviour. It turns out that this requires a much more careful analysis, and we will here restrict to the case where the dynamics are minimal. Our aim is the following statement.

Theorem 3.6. *Suppose that $f|_K$ is minimal and there exists a linear subspace $V \subseteq \mathbb{R}^d$, such that $\rho_K(F) \subseteq v_0 + V^\perp$ for some vector $v_0 \in V$. Further, assume that for any $v \in V$ the deviations parallel to v are unbounded on K . Then there exists a residual set $\mathcal{R} \subseteq K$, such that for all $z \in \mathcal{R}$ there holds*

$$\sup_{n \in \mathbb{N}} D_v(n, z) = -\inf_{n \in \mathbb{N}} D_v(n, z) \quad \forall v \in V.$$

Needless to say that for a minimal irrational pseudo-rotation with unbounded deviations in all directions, as we will construct it below, this implies a rather ‘wild’ behaviour. Our first step is to consider a one-dimensional affine subspace.

Lemma 3.7. *Suppose $f|_K$ is minimal, $\rho_K(F) \subseteq \lambda v + \{v\}^\perp$ and the deviations from the constant rotation parallel to v are not uniformly bounded. Then there exists a residual subset $\mathcal{R} \subseteq K$, such that*

$$\sup_{n \in \mathbb{N}} D_v(n, z) = -\inf_{n \in \mathbb{N}} D_v(n, z) = \infty \quad \forall z \in \mathcal{R}.$$

Proof. Note that there holds

$$\mathcal{R}^+ := \left\{ z \in \mathbb{T}^d \mid \sup_{n \in \mathbb{N}} D_v(n, z) = \infty \right\} = \bigcap_{k \in \mathbb{N}} \mathcal{R}_k^+,$$

where $\mathcal{R}_k^+ := \{z \in \mathbb{T}^d \mid \exists n \in \mathbb{N} : D_v(n, z) > k\}$. The sets $\mathcal{R}_k^+$ are obviously open, and from Lemma 3.1 and the minimality of f it follows that they are also dense (they contain the orbit of the point z_+ in Lemma 3.1). Hence $\mathcal{R}^+$ is residual. The same applies to the set $\mathcal{R}^- := \{z \in \mathbb{T}^d \mid \inf_{n \in \mathbb{N}} D_v(n, z) = -\infty\}$, and the intersection yields the required set $\mathcal{R}$. $\square$

In order to generalise Lemma 3.7 to higher-dimensional hyperplanes, it will be convenient not only to consider the scalar quantities $D_v(n, z)$, but also the directions of the

vectors $D(n, z)$ and their projections to some linear subspace $V \subseteq \mathbb{R}^d$. If $\pi_V : \mathbb{R}^d \rightarrow V$ denotes the orthogonal projection to V , let

$$(3.3) \quad D_V(n, z) := \pi_V(D(n, z)) .$$

As usual, the unit sphere in $\mathbb{R}^d$ is denoted by $\mathbb{S}^{d-1} := \{v \in \mathbb{R}^d \mid \|v\| = 1\}$. Given any $v \in \mathbb{S}^{d-1}$ we let $S^+(v) := \{w \in \mathbb{S}^{d-1} \mid \langle v, w \rangle > 0\}$. We define the set of ‘asymptotic directions’ of $z \in K$ as

$$\mathcal{A}_V(z) := \left\{ v \in \mathbb{S}^{d-1} \mid \forall \varepsilon, r > 0 \exists n \in \mathbb{N} : \|D_V(n, z)\| > r \text{ and } \frac{D_V(n, z)}{\|D_V(n, z)\|} \in B_\varepsilon(v) \right\} ,$$

and let $\mathcal{A}_V^K := \bigcup_{z \in K} \mathcal{A}_V(z)$ and $\mathcal{R}_V(v) := \{z \in \mathbb{T}^d \mid v \in \mathcal{A}_V(z)\}$. The following lemma collects some elementary properties of these sets.

Lemma 3.8. *Suppose $V \subseteq \mathbb{R}^d$ is a linear subspace and $\rho_K(F) \subseteq v_0 + V^\perp$ for some $v_0 \in V$. Then the following holds:*

- (a) *For all $z \in K$, the set $\mathcal{A}_V(z)$ is compact and $\mathcal{A}_V(f(z)) = \mathcal{A}_V(z)$.*
- (b) *If $\sup_{n \in \mathbb{N}} \|D_V(n, z)\| = \infty$, then $\mathcal{A}_V(z)$ is non-empty.*
- (c) *If $\sup_{n \in \mathbb{N}} D_v(n, z) = \infty$ for some $v \in V$, then $\overline{S^+(v)} \cap \mathcal{A}_V(z) \neq \emptyset$.*
- (d) *$\mathcal{R}_V(v)$ is an invariant G_δ -subset of K . In particular, if $\mathcal{R}_V(v)$ contains a point with dense forward orbit, then it is a residual set.*

Proof. (a) The fact that $\mathcal{A}_V(f(z)) = \mathcal{A}_V(z)$ is obvious from the definition. In order to see that $\mathcal{A}_V(z)$ is compact, let

$$A_k(z) := \left\{ v \in \mathbb{S}^{d-1} \mid \exists n \in \mathbb{N} : \|D(n, z)\| > k \text{ and } \frac{D(n, z)}{\|D(n, z)\|} \in B_{1/k}(v) \right\} .$$

Then $\mathcal{A}_V(z) = \bigcap_{k \in \mathbb{N}} \overline{A_k(z)}$ is compact as an intersection of compact sets.

- (b) When the sequence $(D_V(n, z))_{n \in \mathbb{N}}$ is unbounded, then $\overline{A_k(z)} \neq \emptyset \forall k \in \mathbb{N}$, and hence the intersection is non-empty.
- (c) The fact that $\sup_{n \in \mathbb{N}} D_v(n, z) = \infty$ implies that for all $k \in \mathbb{N}$, the sets $\overline{S^+(v)} \cap \overline{A_k(z)}$ are non-empty, and again this carries over to their intersection.
- (d) The invariance of $\mathcal{R}_V(v)$ follows directly from $\mathcal{A}_V(f(z)) = \mathcal{A}_V(z)$. In order to see that $\mathcal{R}_V(v)$ is a G_δ -set, let $R_k(v) := \{z \in \mathbb{T}^d \mid v \in A_k(z)\}$. It is easy to see that these sets are open, and $\mathcal{R}_V(v) = \bigcap_{k \in \mathbb{N}} R_k(v)$. □

Further, we need a piece of linear algebra.

Lemma 3.9. *Suppose $V \subseteq \mathbb{R}^d$ is a linear subspace of dimension $k \leq d$ and $A \subseteq \mathbb{S}^{d-1} \cap V$ is a compact set with the property that*

$$\mathbb{S}^{d-1} \cap V \subseteq \bigcup_{v \in A} \overline{S^+(v)} .$$

Then there exist vectors $v^1, \dots, v^n \in A$ and a linear subspace $\tilde{V} \subseteq V$ of dimension $< k$, such that there holds

$$(\mathbb{S}^{d-1} \cap V) \setminus \bigcup_{i=1}^n S^+(v_i) \subseteq \tilde{V} .$$

Proof. We proceed by induction on the dimension k of V . If $k = 1$, such that $\mathbb{S}^{d-1} \cap V$ consists of exactly two antipodal points, then the statement is obvious. Hence, suppose that it holds for all linear subspaces of dimension $< k$. If

$$\mathbb{S}^{d-1} \cap V \subseteq \bigcup_{v \in A} S^+(v) ,$$

then using compactness we can find $v^1, \dots, v^n \in A$, such that

$$\mathbb{S}^{d-1} \cap V \subseteq \bigcup_{i=1}^n S^+(v^i)$$

and we are finished. (In this case $\dim \tilde{V} = 0$.)

Otherwise, there exists $w \in \mathbb{S}^{d-1} \cap V$, such that $S^+(w) \cap A = \emptyset$. In this case, let $V_0 := V \cap \{w\}^\perp$ and $A_0 := A \cap V_0$. Then there holds

$$\mathbb{S}^{d-1} \cap V_0 \subseteq \bigcup_{v \in A_0} \overline{S^+(v)},$$

since for any $u \in \mathbb{S}^{d-1} \cap V_0$ the set

$$\overline{S^+(u)} \cap A_0 = \bigcap_{\varepsilon > 0} \overline{S^+(w_\varepsilon)} \cap A,$$

where $w_\varepsilon = \frac{w + \varepsilon u}{\|w + \varepsilon u\|}$, is non-empty as the intersection of a nested sequence of non-empty compact sets. Applying the induction assumption to V_0 yields vectors $v^1, \dots, v^n \in A_0$ and a linear subspace $\tilde{V}_0 \subseteq V_0$ of dimension $< k - 1$, such that

$$(\mathbb{S}^{d-1} \cap V_0) \setminus \bigcup_{i=1}^n S^+(v^i) \subseteq \tilde{V}_0.$$

If we now define $\tilde{V}$ as the linear subspace spanned by $\tilde{V}_0$ and w , then the assertions of the lemma are satisfied. $\square$

Proof of Theorem 3.6. We proceed by induction on the dimension k of V . For $k = 1$, the statement is just that of Lemma 3.7. Hence, suppose that it holds for all linear subspaces of dimension $< k$. Since the deviations are unbounded parallel to all vectors $v \in V$, Lemma 3.7 and Lemma 3.8(c) together imply that

$$\mathbb{S}^{d-1} \cap V \subseteq \bigcup_{v \in \mathcal{A}_V^K} \overline{S^+(v)}.$$

Hence $\mathcal{A}_V^K$ satisfies the assumptions of Lemma 3.9, and we obtain vectors $v^1, \dots, v^n$ and a linear subspace $\tilde{V} \subseteq V$, such that

$$\mathbb{S}^{d-1} \cap V \subseteq \tilde{V} \cup \bigcup_{i=1}^n S^+(v^i).$$

By the induction assumption, there exists a residual set $\tilde{\mathcal{R}} \subseteq K$, such that for all $z \in \tilde{\mathcal{R}}$ the deviations parallel to all $v \in \tilde{V}$ are unbounded both above and below. Further, the sets $\mathcal{R}_V(v^1), \dots, \mathcal{R}_V(v^n)$ are residual by Lemma 3.8(d). Hence $\mathcal{R} := \tilde{\mathcal{R}} \cap \mathcal{R}_V(v^1) \cap \dots \cap \mathcal{R}_V(v^n)$ is residual as the intersection of a finite number of residual sets. For any $z \in \mathcal{R}$ and $v \in \mathbb{S}^{d-1} \cap V$ there holds $\sup_{n \in \mathbb{N}} D_v(n, z) = \infty$, since either $v \in \tilde{V}$ or $v \in S^+(v^i)$ for some $i = 1, \dots, n$. Obviously, this also implies $\inf_{n \in \mathbb{N}} D_v(n, z) = -\sup_{n \in \mathbb{N}} D_{-v}(n, z) = -\infty$ for all $v \in V$. Thus $\mathcal{R}$ satisfies the assertions of the theorem. $\square$

As the preceding discussion shows, there is quite a good control about the behaviour of the deviations $D(n, z)$ when these are unbounded and the dynamics are minimal. In contrast to this, the Examples 3.2 show that similar results do not hold in general. The property which was exploited in the construction of these examples was the existence of homotopically non-trivial wandering domains, which is certainly a very strong form of non-recurrence. Hence, an obvious question, which we leave open here, is whether more can be said about systems with intermediate recurrence behaviour (apart from Corollary 3.4). For example, the following would be interesting.

Question 3.10. Suppose that $f \in \text{Homeo}_0(\mathbb{T}^d)$ has lift F with $\rho(F) \subseteq \lambda v + \{v\}^\perp$ and the deviations parallel to v are unbounded.

- (a) If f is non-wandering (does not have any wandering open sets), does this imply that there is some $z \in \mathbb{T}^d$ with $\sup_{n \in \mathbb{N}} |D_v(n, z)| = \infty$?
- (b) If f is topologically transitive, does this imply that there exists some $z \in \mathbb{T}^d$ with $\sup_{n \in \mathbb{N}} D_v(n, z) = \infty$, or even with $\sup_{n \in \mathbb{N}} D_v(n, z) = -\inf_{n \in \mathbb{N}} D_v(n, z) = \infty$?
- (c) Does the statement of Theorem 3.6 remain true if f is non-wandering/topologically transitive?
- (d) If f is topologically transitive, or if f is non-wandering and does not have periodic orbits, does f have sensitive dependence on initial conditions? (In the non-wandering case, the existence of periodic orbits is excluded in order to avoid trivial counter-examples.)

A partial result on question (d) in dimension two will be given in the next section.

Finally, as we have mentioned we will provide some examples with unbounded deviations in all directions.

Proposition 3.11. There exists a minimal pseudo-rotation $f \in \text{Homeo}_0(\mathbb{T}^d)$, such that for any $v \in \mathbb{R} \setminus \{0\}$ there holds $\sup_{n \in \mathbb{N}, z \in \mathbb{T}^d} |D_v(n, z)| = \infty$.

Remark 3.12. The proof below uses the well-known Anosov-Katok-method [13, 14] of fast approximation by rational maps. Using some standard modifications, one may additionally require f to be uniquely ergodic and ρ to be totally irrational. Further, one may require the examples to be smooth, but the rotation vectors will typically be Liouvillean. However, we expect that alternative methods should also yield similar examples with any given irrational rotation vector. The interesting question is whether such examples can also be made smooth, as this is the case for Herman's examples in [11].

Question 3.13. Are there smooth (C^∞ or even real-analytic) irrational pseudo-rotations with Diophantine rotation vector and unbounded mean motion in all directions?

We will carry out the construction only in dimension $d = 2$, again the modifications for the general case are minor. Before we turn to the proof of Proposition 3.11, we need the following auxiliary statement:

Lemma 3.14. Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$ has lift F and $\#\rho(F) = 1$. Further, assume that $u, w \in \mathbb{R}^d$ are linearly independent and for some $z \in \mathbb{R}^2$ there holds

- (i) $\sup_{n \in \mathbb{N}} |D_u(n, z)| = \sup_{n \in \mathbb{Z}} |D_w(n, z)| = \infty$;
- (ii) $\limsup_{n \rightarrow \infty} \frac{|D_u(n, z)|}{|D_w(n, z)| + 1} = \infty$.

Then $\sup_{n \in \mathbb{N}} |D_v(n, z)| = \infty \forall v \in \mathbb{R}^2 \setminus \{0\}$.

Proof. Suppose for a contradiction that for some $v \in \mathbb{R}^2 \setminus \{0\}$ and $C > 0$ there holds $\sup_{n \in \mathbb{N}} |D_v(n, z)| \leq C$. Choose any vector $v' \in \{v\}^\perp \setminus \{0\}$, and write u, w as $u = \lambda_1 v + \lambda_2 v'$ and $w = \mu_1 v + \mu_2 v'$. In order to simplify notation, we will assume $\lambda_1, \lambda_2, \mu_1, \mu_2 \geq 0$. Note that due to property (i), there holds $\lambda_2, \mu_2 \neq 0$. Now, given any $n \in \mathbb{N}$, we have

$$\begin{aligned} \frac{|D_u(n, z)|}{|D_w(n, z)| + 1} &= \frac{|\lambda_1 D_v(n, z) + \lambda_2 D_{v'}(n, z)|}{|\mu_1 D_v(n, z) + \mu_2 D_{v'}(n, z)| + 1} \\ &\leq \frac{\lambda_1 C + \lambda_2 |D_{v'}(n, z)|}{\max\{1, \mu_2 |D_{v'}(n, z)| - \mu_1 C\}} \\ &\leq \max \left\{ \lambda_1 C + \frac{2\mu_1 \lambda_2 C}{\mu_2}, \frac{\lambda_1}{\mu_1} + \frac{2\lambda_2}{\mu_2} \right\}. \end{aligned}$$

(For the last step, it is convenient to distinguished the cases $|D_{v'}(n, z)| < \frac{2\mu_1 C}{\mu_2}$ and $|D_{v'}(n, z)| \geq \frac{2\mu_1 C}{\mu_2}$.) This bound does not depend on n , which clearly contradicts (ii). $\square$

Proof of Proposition 3.11. For the construction, we use the well-known Anosov-Katok-method. Before we start, we need some more notation. Given $f, g \in \text{Homeo}_0(\mathbb{T}^2)$, let $d_{\text{hom}}(f, g) := \sup_{z \in \mathbb{T}^2} \{d(f(z), g(z)), d(f^{-1}(z), g^{-1}(z))\}$. If F is a lift of f , $n \in \mathbb{Z}$ and $z, v, \rho \in \mathbb{R}^d$, let $D_v(f, \rho, n, z) = \langle F^n(z) - z - n\rho, v \rangle$ (hence, we do not suppress f and ρ in the notation as usual, since we will consider sequences of maps and rotation vectors below). Further, by $[x]$ we denote the integer part of $x \in \mathbb{R}$.

By induction, we will choose sequences of rational rotation vectors $\alpha_n = (p_n/q_n, p'_n/q_n)$, toral homeomorphisms h_n and $f_n = h_n \circ R_{\alpha_n} \circ h_n^{-1}$ and integers k_n and K_n with the following properties:

- (i)_n $h_n(0) = 0$;
- (ii)_n $|\alpha_n - \alpha_{n-1}| < 2^{-n}$;
- (iii)_n $d_{\text{hom}}(f_n, f_{n-1}) < 2^{-n}$;
- (iv)_n $\|(F_n^k(z) - z)/k - \alpha_j\| < 2^{-j} \quad \forall j = 1, \dots, n, k = K_j, \dots, 2K_j - 1$;
- (v)_n $|D_{e^1}(f_n, \alpha_n, k_j, 0)| > j \quad \forall j = 2, 4, \dots, 2[\frac{n}{2}]$;
- (vi)_n $|D_{e^2}(f_n, \alpha_n, k_j, 0)| > j \cdot (|D_{e^1}(f_n, \alpha_n, k_j, 0)| + 1) \quad \forall j = 1, 3, \dots, 2[\frac{n+1}{2}] - 1$;
- (vii)_n For all $z \in \mathbb{T}^2$ and $j \leq n$, there holds $\bigcup_{k=0}^{q_j} B_{1/j}(f_n^k(z)) = \mathbb{T}^2$.

Here F_n denotes the lift of f_n . Before we carry out the induction, let us see how (i)–(vii) imply the statement of the proposition. Due to (iii), the limit $f = \lim_{n \rightarrow \infty} f_n \in \text{Homeo}_0(\mathbb{T}^2)$ exists and is well-defined. By (ii), the limit $\alpha = \lim_{n \rightarrow \infty} \alpha_n$ exists, and there holds $|\alpha - \alpha_n| < 2^{-n} \quad \forall n \in \mathbb{N}$. From (iv), we can therefore deduce that for all $k \geq K_j$ and $z \in \mathbb{T}^d$ there holds

$$\|(F^k(z) - z)/k - \alpha\| = \lim_{n \rightarrow \infty} \|(F_n^k(z) - z)/k - \alpha_j\| + 2^{-j} < 2^{-j+1}.$$

(Note that for all $n \geq j$ condition (iv)_n yields $\|(F_n^k(z) - z)/k - \alpha_j\| < 2^{-j} \quad \forall k \geq K_j$.) This implies $\rho(F) = \{\alpha\}$.

From (v), it follows that $D_{e^1}(f, \rho, k_j, 0) = \lim_{n \rightarrow \infty} D_{e^1}(f_n, \alpha_n, k_j, 0) \geq j \quad \forall j \in 2\mathbb{N}$, in particular $\sup_n |D_{e^1}(f, \rho, n, 0)| = \infty$. Similarly, $\sup_n |D_{e^2}(f, \rho, n, 0)| = \infty$ follows from (vi), which also implies

$$\frac{|D_{e^2}(f, \rho, k_j, 0)|}{|D_{e^1}(f, \rho, k_j, 0)| + 1} = \lim_{n \rightarrow \infty} \frac{|D_{e^2}(f_n, \alpha_n, k_j, 0)|}{|D_{e^1}(f_n, \alpha_n, k_j, 0)| + 1} \geq j \quad \forall j \in 2\mathbb{N} + 1.$$

Thus the assumptions of Lemma 3.14 are all satisfied (with $z = 0$, $u = e^2$, and $w = e^1$), and we conclude that f has unbounded deviations in all directions. Finally, the fact that f is minimal follows easily from property (vii).

In order to start the induction, we choose α_0 arbitrary and $h_0 = \text{Id}_{\mathbb{T}^2}$, such that $f_0 = R_{\alpha_0}$. Then (i)₀ is trivial, and (ii)₀–(vii)₀ are still void.

For the induction step $n \rightarrow n+1$, we certainly have to distinguish between odd and even n , due to the asymmetric conditions (v)_n and (vi)_n. However, we will only give the details for even n , since the odd case is similar and even easier. Thus, suppose n is even and $\alpha_0, \dots, \alpha_n, h_0, \dots, h_n, k_0, \dots, k_n$ and $K_0, \dots, K_n$ have been chosen and satisfy (i)_j–(vii)_{j} $\forall j = 0, \dots, n$. Denote the lifts of $h_0, \dots, h_n$ by $H_0, \dots, H_n$.}

Let $C_n^j := \sup_{z \in \mathbb{R}^2, i = \pm 1} \pi_j(H_n^i(z) - z) \quad (j = 1, 2)$, and note that

$$\sup_{k \in \mathbb{Z}, z \in \mathbb{R}^2} |D_{e^1}(f_n, \alpha_n, k, z)| \leq 2C_n^1.$$

Choose a $1/q_n$ -periodic continuous function $\psi : \mathbb{R} \rightarrow \mathbb{R}$, which satisfies $\psi(0) = 0$ and

$$\psi(\theta) \geq (n+1)(2C_n^1 + 1) + C_n^2 \quad \forall \theta \in I,$$

where $I \subseteq (0, 1/q_n)$ is an open and non-empty interval. Let $G(x, y) = (x, y + \psi(x))$ and define h_{n+1} via its lift $H_{n+1} = H_n \circ G$. Then (i)_{n+1} holds, since $G(0) = 0$. Further, as $R_{\alpha_n} \circ G = G \circ R_{\alpha_n}$, there holds $H_{n+1} \circ R_{\alpha_n} \circ H_{n+1}^{-1} = H_n \circ R_{\alpha_n} \circ H_n = F_n$. Due to

the uniform continuity of H_{n+1} and its inverse, we can now ensure that conditions $(ii)_{n+1}$, $(iii)_{n+1}$ and $(v)_{n+1}$ hold just by choosing α_{n+1} sufficiently close to α_n . The same is true for all inequalities in $(iv)_{n+1}$ and $(vi)_{n+1}$, except for the last ones with $j = n + 1$, which do not yet make sense since we have not chosen k_{n+1} and K_{n+1} so far. Similarly, we can require that all but the last equalities in $(vii)_{n+1}$ hold (note that these conditions are open with respect to d_{hom} .)

We will proceed in two steps and choose an intermediate rotation vector $\tilde{\alpha}_{n+1}$ first. Suppose q is large enough, such that $\tilde{\alpha}_{n+1} = (p_n/q_n + 1/qq_n, p'_n/q_n)$ is sufficiently close to α_n in the above sense and $1/q < |I|$. Let $\tilde{f}_{n+1} = H_{n+1} \circ R_{\tilde{\alpha}_{n+1}} \circ H_{n+1}^{-1}$. Choose $k \in \mathbb{N}$, such that $k/q \in I$ and let $k_{n+1} = kq_n$. We claim that

$$(3.4) \quad |D_{e^2}(f_{n+1}, \alpha_{n+1}, k_{n+1}, 0)| > (n+1) (|D_{e^1}(f_{n+1}, \alpha_{n+1}, k_{n+1}, 0)| + 1) .$$

In order to see this, note that the line $L = \mathbb{T}^1 \times \{0\}$ is $R_{\tilde{\alpha}_{n+1}}^{q_n}$ -invariant, and $R_{\tilde{\alpha}_{n+1}}^{k_{n+1}}(0) \in I \times \{0\}$ by the choice of k . Hence, we obtain that

$$(3.5) \quad \begin{aligned} D_{e^2}(\tilde{f}_{n+1}, \tilde{\alpha}_{n+1}, k_{n+1}, 0) &= \pi_2 \left(\tilde{F}_{n+1}^{k_{n+1}}(0) - k_{n+1} \tilde{\alpha}_{n+1} \right) \\ &\stackrel{(i)_{n+1}}{=} \pi_2 \left(H_{n+1} \circ R_{\tilde{\alpha}_{n+1}}^{k_{n+1}}(0) - kp'_n \right) = \pi_2 \left(H_n(k/q, \psi(k/q)) + k(p_n, p'_n) \right) - kp'_n \\ &= \pi_2 \left(H_n(k/q, \psi(k/q)) \right) \geq \psi(k/q) - C_n^2 > (n+1)(2C_n^1 + 1) . \end{aligned}$$

On the other hand, note that $C_{n+1}^1 = C_n^1$, since G leaves the first coordinate invariant. Therefore

$$(3.6) \quad D_{e^1}(\tilde{f}_{n+1}, \tilde{\alpha}_{n+1}, k_{n+1}, 0) \leq 2C_N^1 .$$

Now (3.5) and (3.6) imply (3.4).

In order to complete the induction step, note that due to uniform continuity there exists $\delta > 0$, such that $d(z, z') < \delta$ implies

$$d(H_{n+1}(z), H_{n+1}(z')) < \frac{1}{2(n+1)} .$$

We choose α_{n+1} sufficiently close to $\tilde{\alpha}_{n+1}$, such that all conditions in $(i)_{n+1}$, $(ii)_{n+1}$, $(iii)_{n+1}$, $(v)_{n+1}$ and $(vi)_{n+1}$ are satisfied, as well as all except for the last ones in $(iv)_{n+1}$ and $(vii)_{n+1}$. Further, we require that the orbits of $R_{\alpha_{n+1}}$ are δ -dense in $\mathbb{T}^2$. Then the orbits of $f_{n+1} = H_{n+1} \circ R_{\alpha_{n+1}} \circ H_{n+1}^{-1}$, which are periodic of period q_{n+1} , will be $1/2(n+1)$ -dense in $\mathbb{T}^2$. This ensures the last condition in $(vii)_{n+1}$. Finally, since f_{n+1} is semi-conjugate to $R_{\alpha_{n+1}}$ it has unique rotation vector α_{n+1} , and hence for any suitably large K_{n+1} the last condition in $(iv)_{n+1}$ holds as well. This completes the proof. $\square$

4 Bounded mean motion and Lyapunov stability

Suppose X is a metric space and $f : X \rightarrow X$ is a continuous map. We call a point $z \in X$ ε -Lyapunov stable, if there exists $\delta > 0$, such that

$$(4.1) \quad f^n(B_\delta(z)) \subseteq B_\varepsilon(f^n(z)) \quad \forall n \in \mathbb{N}_0 .$$

This is slightly weaker than the usual notion of a Lyapunov stable point, which requires that for all $\varepsilon > 0$ there exists a $\delta > 0$ with the property (4.1).

f has *sensitive dependence on initial conditions*, if there exists some $\varepsilon > 0$, such that for all $z \in X$ and $\delta > 0$ there exist $z' \in B_\delta(z)$ and $n \in \mathbb{N}_0$, such that $d(f^n(z), f^n(z')) \geq \varepsilon$. In this situation, we call ε a *separation constant* for f . It is easy to see that ε is a separation constant for f if and only if there does not exist an ε -Lyapunov stable point.

Now, assume that $f \in \text{Homeo}_0(\mathbb{T}^2)$ has lift F , and let

$$(4.2) \quad \varepsilon_f := \sup\{\varepsilon > 0 \mid \forall z, z' \in \mathbb{T}^2 : d(z, z') < \varepsilon \Rightarrow d(f(z), f(z')) < 1/2\} .$$

Note that for all $z, z' \in \mathbb{R}^2$ with $d(z, z') < \varepsilon_f$ there holds $d(F(z), F(z')) < \frac{1}{2}$, and therefore $d(F(z), F'(z)) = d(f(\pi(z)), f(\pi(z')))$. In particular, a point $z \in \mathbb{T}^2$ which is an ε_f -Lyapunov stable point for f lifts to an ε_f -Lyapunov stable point for F .

Given any point $z \in \mathbb{R}^2$, we call $\rho(F, z) = \lim_{n \rightarrow \infty} (F^n(z) - z)/n$ the rotation vector of z , whenever the limit exists. We say f is *non-wandering*, if there exists no open set $U \subseteq \mathbb{T}^2$ with $U \cap f^n(U) = \emptyset \ \forall n \in \mathbb{N}$.

Theorem 4.1. *Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$ is non-wandering, and there exists an ε_f -Lyapunov stable point z_0 with totally irrational rotation vector. Then f is an irrational pseudo-rotation with bounded mean motion.*

As an immediate consequence, we obtain

Corollary 4.2. *Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$ is a non-wandering irrational pseudo-rotation with unbounded mean motion. Then f has sensitive dependence on initial conditions.*

We divide the proof of the theorem into the following two lemmas.

Lemma 4.3. *Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$ is non-wandering, and there exists an ε_f -Lyapunov stable point z_0 with totally irrational rotation vector. Then f is an irrational pseudo-rotation.*

Proof. Let F be a lift of f and suppose z_0 is an ε_f -Lyapunov stable point, $\hat{z}_0 \in \pi^{-1}(z_0)$ and

$$(4.3) \quad \lim_{n \rightarrow \infty} (F^n(\hat{z}_0) - \hat{z}_0)/n = \rho \notin \mathbb{Q}^2.$$

As mentioned above, $\hat{z}_0$ is an ε_f -Lyapunov stable point for F . We assume without loss of generality that $\hat{z}_0 = 0$. Choose $\delta > 0$, such that $F^n(B_\delta(0)) \subseteq B_{\varepsilon_f}(F^n(0)) \ \forall n \in \mathbb{N}_0$. Let $U = B_\delta(0)$ and $U_0 = \pi(U)$. Since ρ is not rational, U does not contain any periodic points. The fact that f is non-wandering therefore implies that there exist infinitely many pairs $(m, v) \in \mathbb{N} \times \mathbb{Z}^2$, such that

$$(4.4) \quad F^m(U) \cap (U + v) \neq \emptyset.$$

Given any such pair, let $\tilde{F} := F^m - v$ and $\tilde{\rho} = m\rho - v$, such that

$$(4.5) \quad \lim_{n \rightarrow \infty} \tilde{F}^n(0)/n = \tilde{\rho}.$$

Without loss of generality, we assume $\pi_1(\tilde{\rho}) > 0$. Our aim is to show that $\rho(\tilde{F}) \subseteq \tilde{\rho}\mathbb{R}$, this will then imply the statement of the lemma quite easily. In order to do so, let $V := \bigcup_{n \in \mathbb{N}_0} \tilde{F}^n(U)$. As $\tilde{F}(U) \cap U \neq \emptyset$, the set V is connected. Further, (4.5) together with the Lyapunov stability of $\hat{z} = 0$ implies that for each fixed $k \in \mathbb{N}_0$ there holds

$$(4.6) \quad \lim_{n \rightarrow \infty} \frac{1}{n} d(\tilde{F}^n(U) + (0, k), n\tilde{\rho}) = 0.$$

We claim that for sufficiently large $N \in \mathbb{N}$, the integer translate $V + (0, N)$ is disjoint from V . In order to see this, note that due to (4.6) the sets U and $V + (0, N)$ will be disjoint for sufficiently large N , since only a finite number of iterates of $U + (0, N)$ intersect the strip $[-1, 1] \times \mathbb{R} \supseteq U$. Hence, if the orbit of U intersects $V + (0, N)$, then it must first intersect $U + (0, N)$. However, by the same argument the orbit of U can only intersect a finite number of its vertical integer translates, such that for sufficiently large N we have $V \cap V + (0, N) = \emptyset$ as required.

Now let $a = \inf\{x \in \mathbb{R} \mid (0, x) \in V\}$, $b = \sup\{x \in \mathbb{R} \mid (0, x) \in V + (0, N)\}$ and define W as the union of V , $V + (0, N)$ and the vertical arc from $(0, a)$ to $(0, b)$. Let Y be the unique connected component of $\mathbb{R}^2 \setminus W$ which is unbounded to the left, and $A = \mathbb{R}^2 \setminus Y$. (See Figure 1.) The following two remarks about these objects will be helpful below:

First, from (4.6) we can deduce for any $m_1 < \rho_2/\rho_1 < m_2$ there exist constants c_1, c_2 , such that

$$(4.7) \quad A \subseteq \{(x, y) \in \mathbb{R}^2 \mid m_1 x + c_1 \leq y \leq m_2 x + c_2\}.$$




Figure 1: The construction of the sets V and W on the right, and the set A on the left.

Secondly, due to the definition of W , its connectedness and the fact that it ‘stretches out to infinity’ by (4.6), the set $S := (\mathbb{R}^+ \times \mathbb{R}) \setminus A = (\mathbb{R}^+ \times \mathbb{R}) \cap Y$ consists of exactly two connected components. These can be defined as follows: Fix any $\zeta_0 \in Y$ with $\pi_1(\zeta) < 0$. For any $\zeta \in S$, there is a path γ_ζ from ζ to ζ_0 . Let x_ζ be the second coordinate of the first point in which γ_ζ intersects the vertical axis. The fact whether x_ζ lies above b or below a does not depend on the choice of the path, since this would contradict the connectedness of W . It is now easy to see that $S^- = \{\zeta \in S \mid x_\zeta > b\}$ and $S^+ = \{\zeta \in S \mid x_\zeta < a\}$ form a partition of S into two connected components.

Claim 4.4. *There exists some constant $\alpha > 0$, such that $z \in A$ and $\pi_1(z) \geq \alpha$ imply $\tilde{F}(z) \in A$ and $\tilde{F}^{-1}(z) \in A$*

Proof. We show that there exists a constant $\alpha' > 0$, such that $\pi_1(z) \geq \alpha'$ and $z \in S$ implies $\tilde{F}(z) \in S$ and $\tilde{F}^{-1}(z) \in S$. If we let

$$(4.8) \quad M := \sup_{z \in \mathbb{R}^2} \|\tilde{F}(z) - z\| = \sup_{n \in \mathbb{R}^2} \|\tilde{F}^{-1}(z) - z\| ,$$

then for any $z \in \pi_1^{-1}[\alpha' + M, \infty)$ this implies that there holds $\tilde{F}(z) \in S \Rightarrow z \in S$, and hence $z \in A \Rightarrow \tilde{F}(z) \in A$. The same applies to $\tilde{F}^{-1}(z)$. Thus we can choose $\alpha = \alpha' + M$.

From (4.6) we deduce that there exists some $n_0 \in \mathbb{N}$, such that

$$\pi_1 \circ \tilde{F}^n(U) \subseteq (4M, \infty) \quad \forall n \geq n_0 .$$

Choose $\alpha' > 0$, such that for all $j \leq n_0$ there holds $\pi_1 \circ \tilde{F}^j(U) \subseteq [0, \alpha')$. We obtain the following statement:

$$(4.9) \quad \pi_1 \circ \tilde{F}^n(U) \cap [\alpha', \infty) \neq \emptyset \quad \Rightarrow \quad \pi_1 \circ \tilde{F}^k(U) \cap [0, 4M] = \emptyset \quad \forall k \geq n .$$

Of course, the same statement applies to $U + (0, N)$.

Due to (4.7), it is possible to choose some $K > 0$, such that

$$B := [0, 4M] \times [K - M, \infty) \subseteq S^+ .$$

Let $z^* = (3M, K)$ and fix $z \in S^+$ with $\pi_1(z) \geq \alpha'$. Then, since S^+ is open and connected, there exists a simple path $\gamma : [0, 1] \rightarrow S^+$ from z to z^* . We claim that γ can be chosen such that its image contained in $S^+ \cap \pi_1^{-1}[3M, \infty)$.

Suppose not, and let $t_0 := \min\{t \in [0, 1] \mid \gamma(t) \in B\}$ and $\Gamma = \{\gamma(t) \mid t \in [0, t_0]\}$. Then Γ divides the set $([0, \pi_1(z)] \times \mathbb{R}) \setminus B$ into exactly two connected components D^+ and D^- , which are unbounded above, respectively below (see Figure 2). Now, if $\bar{W}$ does not intersect $D^+ \cap \pi_1^{-1}[0, 4M)$, then $D^+ \cap \pi_1^{-1}[0, 4M) \subseteq S^+$, and it is easy to see that in this case either γ does not intersect $\pi_1^{-1}[0, 3M)$, or we can modify it to that end. Otherwise, there must




Figure 2: The path γ divides $([0, \pi_1(z)] \times \mathbb{R}) \setminus B$ into two connected components D^+ and D^- . As argued in the proof, the set W (and hence A) cannot ‘reach back’ into D^+ too far.

be some $k \in \mathbb{N}$, such that $F^k(U)$ or $F^k(U) + (0, N)$ intersects $D^+ \cap \pi_1^{-1}[0, 4M)$. However, since U intersects D^- and the set $\bigcup_{i=0}^k \tilde{F}^i(U)$ is connected, this implies that there must be some $n \leq k$, such that $\pi_1 \circ \tilde{F}^n(U)$ intersects $[\pi_1(z), \infty] \subseteq [\alpha', \infty]$. This contradicts (4.9).

Summarising, we have found a path γ from z to z^* , which is contained in $S^+ \cap \pi_1^{-1}[3M, \infty)$. In particular, γ is contained in the complement of $\overline{W}$. Consequently, the path $\tilde{F} \circ \gamma$ is contained in the complement of $\tilde{F}(\overline{W})$. At the same time, it is also contained in $\pi_1^{-1}[2M, \infty)$. However, it follows from the construction of W and the definition of M that $\tilde{F}(\overline{W}) \cap \pi_1^{-1}[2M, \infty) = \overline{W} \cap \pi_1^{-1}[2M, \infty)$. Hence, the path $\tilde{F} \circ \gamma$ is also contained in the complement of $\overline{W}$. Furthermore, it joins $\tilde{F}(z)$ to the point $\tilde{F}(z^*)$, which is contained in $B \subseteq S^+$. This implies that $\tilde{F}(z)$ is equally contained in S^+ . When $z \in S^-$, the argument is similar. In the same way, one can show that $\tilde{F}^{-1}(z) \in S$, which completes the proof.

○

Using this ‘almost-invariance’ of A , we can now prove the following.

Claim 4.5. *There holds $\rho(\tilde{F}) \subseteq \tilde{\rho}\mathbb{R}$.*

Proof. Suppose for a contradiction that $\sigma \in \rho(\tilde{F}) \setminus \tilde{\rho}\mathbb{R}$. Without loss of generality, we may assume that σ is an extremal point of $\rho(\tilde{F})$. In this case, σ is realised by some orbit [2], that is, there exists some $z_1 \in \mathbb{R}^2$ with $\lim_{|n| \rightarrow \infty} (\tilde{F}^n(z_1) - z_1)/n = \sigma$. Note that the limit is take two-sided.

There holds $\sigma \neq 0$. Hence, by permuting the coordinates if necessary, we may assume $\pi_1(\sigma) \neq 0$. We first assume that $\pi_1(\sigma) > 0$. In this case, there exists some $r \in \mathbb{R}$, such that $\pi_1(\tilde{F}^n(z_1) - z_1) > -r \forall n \in \mathbb{N}_0$. Now, there holds $\mathbb{R}^+ \times \mathbb{R} \subseteq \bigcup_{j \in \mathbb{N}} A + (0, jN)$. Replacing z_1 by an integer translate if necessary, we may therefore assume $z_1 \in A$ and $\pi_1(z_1) \geq \alpha + r$. In this case, Claim 4.4 implies $\tilde{F}^n(z_1) \in A \forall n \in \mathbb{N}_0$, and therefore (4.7) yields $\sigma \in \tilde{\rho}\mathbb{R}$, contradicting our assumption. When $\pi_1(\sigma) < 0$, we can proceed in the same way by considering the backwards orbit of z_1 .

○

Now we can conclude the proof of the lemma. Due to (4.3), and using the fact that ρ is totally irrational and v is an integer vector, there exists some $n_0 \in \mathbb{N}$, such that for all $n \geq n_0$ there holds $d(F^n(z) - z, \mathbb{R}v) > 2\varepsilon_f$. It follows that $F^n(U) \cap (U + \lambda v) = \emptyset \forall n \geq n_0, \lambda \in \mathbb{R}$. Hence, we can find a second pair (m', v') that satisfies (4.4), such that the

vectors v and v' are linearly independent. Now, applied to (m, v) , Claim 4.4 yields

$$(4.10) \quad \rho(F) = \frac{\rho(\tilde{F})}{m} + \frac{v}{m} \subseteq \left(\rho - \frac{v}{m} \right) \mathbb{R} + \frac{v}{m}.$$

Similarly, we obtain

$$(4.11) \quad \rho(F) \subseteq \left(\rho - \frac{v'}{m'} \right) \mathbb{R} + \frac{v'}{m'}$$

for the pair (m', v') . As ρ is totally irrational and the rational vectors v/m and v'/m' are linearly independent, the vectors $\rho - v/m$ and $\rho - v'/m'$ are linearly independent as well. Thus (4.10) and (4.11) together imply that $\rho(F)$ is reduced to a singleton. Since the rotation vector ρ is realised on the orbit of z_0 , we must have $\rho(F) = \{\rho\}$. $\square$

Lemma 4.6. *Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$ is a non-wandering irrational pseudo-rotation, and there exists an ε_f -Lyapunov stable point z_0 . Then f has bounded mean motion.*

Proof. We use the same definitions and notation as in the proof of the preceding lemma. As in the proof of Proposition 3.11, we write $D_v(f, \rho, n, z)$ for the deviations from the constant rotation, in order to emphasize the dependence both on the map and on the rotation vector. First, let u be orthogonal to $\rho - v/m$ with norm $\|u\| = 1$, where the pair (v, m) is chosen as in (4.4). Suppose that the deviations $D_u(f, \rho, n, z_0)$ are not uniformly bounded in $n \in \mathbb{N}_0$, without loss of generality

$$(4.12) \quad \sup_{n \in \mathbb{N}_0} D_u(f, \rho, n, z_0) = \infty.$$

In this case, an easy computation yields that the deviations $D_u(\tilde{f}, \tilde{\rho}, n, z_0)$ are equally unbounded in $n \in \mathbb{N}_0$. (By $\tilde{f}$ we denote the total homeomorphism induced by $\tilde{F}$.) We want to lead this to a contradiction.

Assume without loss of generality that $\pi_1(\tilde{\rho}) > 0$. Note that since $\tilde{f}$ is an irrational pseudo-rotation, the quantities $(\tilde{F}^n(z) - z)/n$ converge to $\tilde{\rho}$ uniformly on $\mathbb{R}^2$. It follows that there exists some $n_0 \in \mathbb{N}$, such that there holds

$$(4.13) \quad \pi_1(\tilde{F}^n(z) - z) \geq M + 1 \quad \forall n \geq n_0, z \in \mathbb{R}^2,$$

where M is defined as in (4.8).

Together with the Lyapunov stability of z_0 and its lift, the fact that the deviations $D_u(\tilde{f}, \tilde{\rho}, n, z_0)$ are unbounded implies that for every $r > 0$ there exists some $n \in \mathbb{N}_0$, such that

$$\langle z, u \rangle \geq r \quad \forall z \in \tilde{F}^n(U).$$

It follows that for all $k \in \{n - n_0, \dots, n + n_0\}$ and all $z \in \tilde{F}^k(U)$ there holds $\langle z, u \rangle \geq r - n_0 M$. By (4.13), we obtain that there exists an interval I of length $2M$, such that all $z \in A \cap \pi_1^{-1}(I)$ satisfy $\langle z, u \rangle \geq r - n_0 M - N$. Furthermore, we may assume that I lies to the right of the point $n_0 M + 1$.

Now, as argued in the proof of Lemma 4.3, any point in $\mathbb{T}^2$ has a lift in the compact set $A_0 := A \cap \pi_1^{-1}(n_0 M, n_0 M + 1]$. Furthermore, if n_0 is chosen sufficiently large, such that $n_0 M \geq \alpha$ with α as in Claim 4.4, then the forward orbit of this lift under $\tilde{F}$ will stay in A all the time. However, any such orbit has to pass through the set $\pi_1^{-1}(I)$. By choosing $r \geq n_0 M + N + \sup_{z \in A_0} \|z\| + 1$, we obtain that for every $z \in \mathbb{T}^2$ there holds $\sup_{n \in \mathbb{N}} D_u(\tilde{f}, \tilde{\rho}, n, z) \geq 1$. This contradicts Lemma 3.1.

Hence, we may assume that z_0 has bounded deviations parallel to u . It follows that that the set A lies within a bounded distance of the semi-line $\tilde{\rho}\mathbb{R}$. Since every point in $\mathbb{T}^2$ has a lift whose orbit stays forever in A , this further implies that the deviations $D_u(\tilde{f}, \tilde{\rho}, n, z)$ are uniformly bounded in n and z . Consequently, the same holds for the deviations $D_u(f, \rho, n, z)$.

Now replace the pair (m, v) by (m', v') with v and v' linearly independent, as at the end of the proof of Lemma. Then we obtain another, linearly independent vector u' , such

that the deviations $D_{u'}(f, \rho, n, z)$ are uniformly bounded. This implies that the vectors $D(f, \rho, n, z)$ are uniformly bounded as well. $\square$

Remark 4.7. *The only time that we used the fact that the rotation vector ρ of z_0 is totally irrational, and not only non-rational, in the above proof was to deduce that there exist two pairs (m, v) and (m', v') satisfying (4.4), with linearly independent vectors v and v' . Hence, if ρ is neither rational nor totally irrational, the above construction still yields the following statement:*

Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$ is non-wandering, and there exists an ε_f -Lyapunov stable point z_0 with a rotation vector ρ that is neither rational nor totally irrational. Then there exists a vector $w \in \mathbb{Q}^2$, such that $\rho(f) \subseteq \rho + (\rho - w)\mathbb{R}$ and the deviations orthogonal to $\rho - w$ are uniformly bounded.

References

- [1] T. Jäger. Linearisation of conservative toral homeomorphisms. Preprint, 2008.
- [2] M. Misiurewicz and K. Ziemian. Rotation sets for maps of tori. *Journal of the London Mathematical Society*, 40:490–506, 1989.
- [3] J. Franks. Realizing rotation vectors for torus homeomorphisms. *Transactions of the American Mathematical Society*, 311(1):107–115, 1989.
- [4] J. Llibre and R.S. MacKay. Rotation vectors and entropy for homeomorphisms of the torus isotopic to the identity. *Ergodic Theory and Dynamical Systems*, 11:115–128, 1991.
- [5] M. Misiurewicz and K. Ziemian. Rotation sets and ergodic measures for torus homeomorphisms. *Fundamenta Mathematicae*, 137(1):45–52, 1991.
- [6] J. Stark, U. Feudel, P.A. Glendinning, and A. Pikovsky. Rotation numbers for quasiperiodically forced monotone circle maps. *Dynamical Systems*, 17(1):1–28, 2002.
- [7] T. Jäger and J. Stark. Towards a classification for quasiperiodically forced circle homeomorphisms. *Journal of the LMS*, 73(3):727–744, 2006.
- [8] K. Bjerklov and T. Jäger. Rotation numbers for quasiperiodically forced circle maps – Mode-locking vs strict monotonicity. Preprint, 2006.
- [9] Y Yi. On almost automorphic oscillations. *Fields Inst. Commun.*, 42:75–99, 2004.
- [10] W. Huang and Y. Yi. Almost periodically forced circle flows. Preprint, 2007.
- [11] M. Herman. Une méthode pour minorer les exposants de Lyapunov et quelques exemples montrant le caractère local d’un théorème d’Arnold et de Moser sur le tore de dimension 2. *Commentarii Mathematici Helvetici*, 58:453–502, 1983.
- [12] B. Fayad. Weak mixing for reparameterized linear flows on the torus. *Ergodic Theory and Dynamical Systems*, 22(1):187–201, 2002.
- [13] D. Anosov and A. Katok. New examples in smooth ergodic theory. ergodic diffeomorphisms. *Transactions of the Moscow Mathematical Society*, 23:1–35, 1970.
- [14] B. Fayad and A. Katok. Constructions in elliptic dynamics. *Ergodic Theory and Dynamical Systems*, 24(5):1477–1520, 2004.